

A note on the genus of certain curves over finite fields

Rainer Furhmann

Fernando Torres*

Abstract

We prove the following result which was conjectured by Stichtenoth and Xing: let g be the genus of a projective, irreducible non-singular curve over the finite field $\mathbb{F}_{q^2}$ and whose number of $\mathbb{F}_{q^2}$ -rational points attains the Hasse-Weil bound; then either $4g \leq (q-1)^2$ or $2g = (q-1)q$.

1. Throughout this note by a curve we mean a projective, irreducible non-singular algebraic curve defined over the finite field $k := \mathbb{F}_{q^2}$ with q^2 elements. This note is concerning with the genus g of a curve X whose number of k -rational points $\#X(k)$ attains the Hasse-Weil bound, that is,

$$\#X(k) = q^2 + 2gq + 1. \quad (1)$$

These curves are very useful for applications to coding theory (see e.g. [Sti], [Tsf-Vla]). It is known that $2g \leq (q-1)q$ ([Sti, V.3.3]), and that the Hermitian curve is the only one that satisfies (1) and whose genus is $(q-1)q/2$ [R-Sti]. Here we prove the following result which was conjectured by Stichtenoth and Xing in [Sti-X].

Theorem. Let X be a curve of genus g , and suppose that X satisfies (1). Then

$$4g \leq (q-1)^2 \quad \text{or} \quad 2g = (q-1)q.$$

We prove this theorem by using [Sti-X, Prop. 1], Rück and Stichtenoth's [R-Sti, Lemma 1] and by using a particular case of the theory of Frobenius orders sequence associated to linear system of curves which was developed by Stöhr and Voloch [S-V] as a way of improving the Hasse-Weil bound. We also state another proof of Rück and Stichtenoth's result (cf. [R-Sti]) concerning Hermitian curves (see §4). We remark that Hirschfeld, Storme, Thas and Voloch also stated a characterization of Hermitian curves by using some results from [S-V] (see [HSTV]). In §3 we use ideas from the proof of [HSTV, Lemma 1].

In [F-T] is considered the case of curves satisfying (1) and whose genus is bounded from above by $(q-1)^2/4$.

We are indebted to Professor J.F. Voloch for pointing out to us that the proof of the theorem in the previous version of this paper was incomplete.

*Supported by a grant from the International Atomic Energy Agency and UNESCO.

2. Let X be a curve of genus g , satisfying the hypothesis (1). The starting point of the proof is the fact that there exists a k -rational point $P_0 \in X$ such that q and $q + 1$ are not-gaps at P_0 [Sti-X, Prop. 1]. Then the linear system

$$\mathcal{D} = g_{q+1}^{r+1} := |(q+1)P_0|$$

is simple. Thus the genus g of X and the dimension $r + 1$ of $\mathcal{D}$ are related by Castelnuovo's genus bound for curves in projective spaces ([C], [ACGH, p. 116], [Ra, Corollary 2.8]). Thus we have

$$2g \leq M(q - r + e), \quad (2)$$

where M is the biggest integer $\leq q/r$ and $e := q - Mr$.

Claim. 1 ([Sti-X, Prop. 3]) *If $r + 1 \geq 3$, then $4g \leq (q - 1)^2$.*

Proof. From (2) we have

$$2g \leq \frac{(q - e)(q - r + e)}{r} \leq \frac{(2q - r)^2}{4r}.$$

This implies the result. $\square$

From now on we consider $r + 1 = 2$. We apply [S-V] to the linear system $\mathcal{D} = g_{q+1}^2 = |(q+1)P_0|$. For $P \in X$ let $0 = j_0(P) < j_1(P) < j_2(P) \leq q + 1$ denote the $(\mathcal{D}, P)$ -orders ([S-V, §1]. The fact that q and $q + 1$ are non-gaps at P_0 implies

$$j_1(P_0) = 1 \quad \text{and} \quad j_2(P_0) = q + 1.$$

To compute $j_2(P)$ for $P \in X(k)$ we use the fact that the linear system $|(q+1)P|$ is equivalent to $\mathcal{D}$ ([R-Sti, Lemma 1]). Then

$$j_2(P) = q + 1, \quad (3)$$

provided $P \in X(k)$.

Let denote by $0 = \nu_0 < \nu_1$ the k -Frobenius order sequence of $\mathcal{D}$ (see [S-V, Prop. 2.1]). ν_1 satisfies

$$\nu_1 \leq j_2(P) - j_1(P) = q + 1 - j_1(P), \quad (4)$$

for each $P \in X(k)$ [S-V, Corollary 2.6].

Associated to $\mathcal{D}$ there exists a positive divisor S of degree

$$\deg(S) = \nu_1(2g - 2) + (q^2 + 2)(q + 1) \quad (5)$$

whose support contains $X(k)$. Moreover by [S-V, Prop. 2.4] we have

$$v_P(S) \geq j_1(P) + (j_2(P) - \nu_1) \geq q + 2 - \nu_1. \quad (6)$$

Claim. 2 *Suppose $4g > (q - 1)^2$. Then*

- (i) $\nu_1 = q$.
- (ii) $j_1(Q) = 1$ for $Q \in X(k)$.

Proof. Statement (ii) follows from statement (i) and relation (4). Applying (4) for P_0 we also obtain $\nu_1 \leq q$. Now by using relations (6), (1) and (5) we get

$$(q-1)(\nu_1(q+1)-q) \geq 2g(q^2 - \nu_1(q+1) + 2q).$$

Finally by using the hypothesis on g we get $\nu_1 \geq q$ and we are done. $\square$

3. Proof of the Theorem. Let assume that

$$4g > (q-1)^2.$$

From the fact that q and $q+1$ are non-gaps at P_0 we have $2g \leq (q-1)q$ (see e.g Jenkins's [J]).

Let R be the divisor on X supporting the $\mathcal{D}$ -Weierstrass points, and let denote by $0 = \epsilon_0 < \epsilon_1 < \epsilon_2$ the order sequence of $\mathcal{D}$ (see [S-V, §1]). The degree of R is given by

$$\deg(R) = (\epsilon_1 + \epsilon_2)(2g - 2) + 3(q + 1).$$

For each $P \in X$ we have $v_P(R) \geq \sum_{i=0}^2 (j_i(P) - \epsilon_i)$. Now since $\epsilon_i \leq j_i(P)$ for each i , from Claim 2 we have $\epsilon_1 = 1$. Moreover by [S-V, Prop. 2.1] we have $\epsilon_2 = \nu_1$. Thus $\epsilon_2 = q$ by Claim 2. Hence by relation (3) we have

$$v_P(R) \geq 1$$

for each $P \in X(k)$. Consequently by (1) and $\deg(R)$ it follows that $2g \geq (q-1)q$, which proves the result. $\square$

4. Remark. We close this note by proving that a curve satisfying (1) and whose genus is $(q-1)q/2$ is k -isomorphic to the so called Hermitian curve which is defined by an equation of type

$$y^q + y = x^{q+1}.$$

This result was proved in [R-Sti]. Here we give a different proof which is inspired on the example stated in [S-V, p. 16].

Let P_0 be as in §2 and let $x, y \in k(X)$ such that

$$\operatorname{div}_\infty(x) = qP_0 \quad \text{and} \quad \operatorname{div}_\infty(y) = (q+1)P_0.$$

Then by the Riemann-Roch theorem the k -dimension of the linear system $|q(q+1)P_0|$ is equal to $(q+1)(q+2)/2 - 1$. Since

$$\#\{(i, j) \in \mathbb{N} \times \mathbb{N} : iq + j(q+1) \leq q(q+1)\} = \frac{(q+1)(q+2)}{2} + 1,$$

there exists a non-trivial k -linear relation:

$$F = \sum_{iq+j(q+1) \leq q(q+1)} a_{i,j} x^i y^j = 0, \tag{7}$$

where $a_{q+1,0} \neq 0$ and $a_{0,q} \neq 0$. X is isomorphic to the plane curve defined by F . After a k -projective transformation we may assume $a_{q+1,0} = a_{0,q} = 1$ and $a_{q,0} = 0$. The fact that $\nu_1 = q > 1$ implies

$$(x - x^{q^2})F_x = (y^{q^2} - y)F_y, \tag{8}$$

where F_x (resp. F_y) stands for the partial derivative with respect to the variable x (resp. y) of F . Let v denote the valuation associated to P_0 . From (8) we find that $F_y \neq 0$ and $v(F_y) = 0$. Thus $F_y = a_{0,1}$. Now we compute y^{q^2} by means of (7) and hence from (8) we get $a_{0,1} = 1$, $a_{i,j} = 0$ for $(i,j) \notin \{(0,0), (0,q), (q+1,0)\}$. Finally by another k -projective transformation we may assume $a_{0,0} = 0$ and we find that F defines the Hermitian curve.

References

- [ACGH] Arbarello, E.; Cornalba, M.; Griffiths, P.A. and Harris, J.: Geometry of algebraic curves, Vol. I, Springer-Verlag, New York 1985.
- [C] Castelnuovo, G.: Ricerche di geometria sulle curve algebriche, Atti. R. Acad. Sci. Torino **24**, 196–223 (1889).
- [F-T] Furhmann, R.; Torres, F.: Curves over finite fields with maximal number of rational points, (In preparation).
- [HSTV] Hirschfeld, J.W.P.; Storme, L.; Thas, J.A. and Voloch, J.F.: A characterization of Hermitian curves, J. Geom. **41**, 72–78 (1991).
- [J] Jenkins, J.A.: Some remarks on Weierstrass points, Proc. Amer. Math. Soc. **44**, 121–122 (1974).
- [Ra] Rathmann, J.: The uniform position principle for curves in characteristic p , Math. Ann. **276**, 565–579 (1987).
- [R-Sti] Rück, H.G.; Stichtenoth, H.: A characterization of Hermitian function fields over finite fields, J. Reine. Angew. Math. **457**, 185–188 (1994).
- [Sti] Stichtenoth, H.: Algebraic functions fields and codes, Springer-Verlag, Berlin, 1993.
- [Sti-X] Stichtenoth, H.; Xing, C.: The genus of maximal functions fields, Manuscripta Math. **86**, 217–224 (1995).
- [S-V] Stöhr, K.O.; Voloch, J.F.: Weierstrass points and curves over finite fields, Proc. London Math. Soc. (3) **52**, 1–19 (1986).
- [Tsf-Vla] Tsfasman, M.; Vladut, S.G.: Algebraic-Geometric Codes, Kluwer Academic Publishers, Dordrecht-Boston-London 1991.